

JOURNAL OF DIFFERENTIAL EQUATIONS 6, 314–325 (1969)

Singular Perturbation of Linear Partial  
Differential Equation

H. S. NUR

*Department of Mathematics, Fresno State College, Fresno, California 93726*

Received June 28, 1968

## I

The aim of this paper is to show the convergence of certain solution of the homogeneous linear partial differential equation

$$L_\epsilon(u) = \epsilon \delta_t^2 u + P_1(\delta_x, t) \delta_t u + P_2(\delta_x, t, \epsilon) u = 0 \quad (1)$$

as  $\epsilon > 0$  goes to zero<sup>1</sup> to a solution of the limit equation ( $\epsilon = 0$ )

$$L_0(u) = 0 \quad (1^0)$$

where  $P_i(\delta_x, t, \epsilon)$  ( $i = 1, 2$ ) are polynomials in  $\delta_{x_1}, \delta_{x_2}, \dots, \delta_{x_m}$  and  $\epsilon$  with complex coefficients which are  $C^\infty$  functions in the real parameter  $t$  in  $J = [0, T]$  for some fixed positive number  $T$ .

We demand here that  $P_1(\delta_x, t)$  does not depend on  $\epsilon$ .

This relationship has been discussed in [1] for equation of order  $n$  but with constant coefficients. Here we permit the coefficients to depend on the real parameter  $t$ .

[Here I wish to thank the referee for his suggestion which enables me to reduce the size of this paper considerably.]

**DEFINITION (1).** For any integer  $p > 0$  and any function  $u(x)$  in  $C_0^\infty$  let the norm of  $u$  be defined as follows:

$$\|u\|_p^2 = \int_{E^m} |(1 + |z|^2)^{p/2} \hat{u}(z)|^2 dz \quad (2)$$

<sup>1</sup> In what follows there will be for each  $\epsilon$  certain exceptional sets of  $z$  ( $z$  in  $E^m$ ) of measure zero for which our conclusions do not apply. In order to be able to draw inferences as  $\epsilon \rightarrow 0$ ,  $\epsilon > 0$  we wish to be able to disregard these sets.

Now let the notion  $\epsilon \rightarrow 0$  be henceforth interpreted as meaning “ $\epsilon$  tends to zero through an arbitrary sequence of positive numbers.” Then all of the corresponding exceptional sets will still be according to Theorem (2) in [1] a countable union of sets of measure zero and accordingly itself has measure zero.

( $\wedge$  is the Fourier transformation). The space  $C_0^\infty$  with the norm (3) gives a Hilbert space. We shall call it  $H_{p,x}$  space.

DEFINITION (2). Let  $u(t)$  be a variable element of  $H_{p,x}$  depending a real parameter  $t$  in  $J$ . Then  $u(t)$  is  $H_{p,x}$  continuous in  $t$  if the mapping  $t$  in  $J \rightarrow u(t)$  in  $H_{p,x}$  is continuous, i.e.  $t \rightarrow t_0$  in  $J$  implies that  $u(t) \rightarrow u(t_0)$  in  $H_{p,x}$ . We say that  $u(t)$  is  $H_{p,x}$ -differentiable at  $t = t_0$ , if there exists a function  $g(t)$  in  $H_{p,x}$  such that  $(t - t_0)^{-1}[u(t) - u(t_0)] \rightarrow g(t_0)$  in  $H_{p,x}$  as  $t \rightarrow t_0$ . Then we denote  $g(t)$  by  $(d/dt)u(t)$ .

DEFINITION (3). We say that equation (1) is  $H_{p,x}$ -stable equation for  $\epsilon \rightarrow 0$  in  $0 \leq t \leq T$  with respect to a particular solution  $u(t)$  of (1<sup>0</sup>) for ( $\epsilon = 0$ ), if and only if,  $u_\epsilon(t) \rightarrow u_0(t)$  in  $H_{p,x}$  for  $t$  in  $J$  whenever  $u_\epsilon(t, x)$  is an  $H_{p,x}$  solution of equation (1) with the property:

$$u_\epsilon(0) \rightarrow u_0(0) \quad (3)$$

in  $H_{p,x}$  and there exists a function  $F(x)$  in  $H_{p,x}$  with the following property:

$$|\delta_\epsilon u(0, z)| \leq |\hat{F}(z)| \quad (4)$$

for all small  $\epsilon > 0$ .

## II

In this section we shall derive an asymptotic expansion for  $Y_j(t, z, \epsilon)$  (the solution of equation (2)) with the initial conditions

$$\delta_\epsilon^{k-1} Y_j(0, z, \epsilon) = \delta_{jk}. \quad (5)$$

Just for the simplicity, we made the order of equation (1) be 2. Otherwise it would be harder to compute the constants in  $Y_j(t, z, \epsilon)$  by using the above initial conditions.

As in [1] we associate equation (1) with the linear ordinary differential equation

$$\epsilon D_t^2 u + P_1(iz, t) D_t u + P_2(iz, t, \epsilon) u = 0 \quad (6)$$

for  $z$  in  $E^m$ .

LEMMA (1). Let  $Y_j(t, z, \epsilon)$  be the solutions of equation (6) with the initial conditions (5) satisfying the following condition: There exist two constants  $\epsilon_0 > 0$  and  $C$  such that

$$\text{Sup } |Y_j(t, z, \epsilon)| \leq C \quad (7)$$

for all  $t$  in  $J$  and  $0 < \epsilon \leq \epsilon_0$  and the sup is taken over all  $z$  in  $E^m$ . Then real  $[P_1(iz, t)]$  is nonnegative for all  $t$  in  $J$ .

*Proof.* We rewrite equation (6) as

$$D[\epsilon Dy + P_1(iz, t)y] + \tilde{P}_2(iz, t, \epsilon)y = 0$$

where;  $\tilde{P}_2 = (dP_1/dt) + P_2$ . Finding  $u$  from the equation

$$D_t u + \tilde{P}_2(iz, t, \epsilon)y = 0$$

and substituting it in the equation

$$\epsilon D_t y + P_1(iz, t)y = u$$

we find that

$$y = \exp \left[ - (1/\epsilon) \int P_1(iz, t) dt \right] \\ \times \left[ \int^t \exp \left[ (1/\epsilon) \int^s P_1(iz, t) dt \right] \frac{1}{\epsilon} \left( \int^s - \tilde{P}_2(iz, t, \epsilon) y dt + C_2 \right) ds + C_1 \right]$$

For  $y = Y_1(iz, t, \epsilon)$  using the initial conditions (5) we find  $C_2 = 0$  and  $C_1 = 1$ . Consequently

$$Y_1(iz, t, \epsilon) = [1 + G(z, t, \epsilon)] \exp \left[ - (1/\epsilon) \int P_1(iz, t) dt \right]$$

where

$$G(z, t, \epsilon) = \int^t \exp \left[ (1/\epsilon) \int^s P_1(iz, t) dt \right] \frac{1}{\epsilon} \left( \int^s - \tilde{P}_2(iz, t, \epsilon) Y_1 dt \right) ds.$$

Using the triangular inequality we get

$$|Y_1(t, z, \epsilon)| \geq \left| \exp \left[ -\frac{1}{\epsilon} \int P_1(iz, t) dt \right] \right| |1 - |G(z, t)||. \quad (*)$$

Notice that  $Y_1(t, z, \epsilon)$  is bounded and  $\tilde{P}_2(iz, t, \epsilon)$  is a polynomial in  $\epsilon$ , therefore if, real  $P_1(iz, t) < 0$  then Ascoli's theorem implies that

$$|G(z, t, \epsilon)| \rightarrow 0$$

as  $\epsilon \rightarrow 0$ . The inequality (\*) shows that

$$|Y_1(t, z, \epsilon)| \rightarrow \infty$$

as  $\epsilon \rightarrow 0$  and this is a contradiction. The lemma is proved.

Let  $u^{[k]} = \epsilon^k d^k u / dt^k$ . Inserting this in equation (6), we have

$$Ay \equiv y^{[2]} + P_1(iz, t) y^{[1]} + \epsilon P_2(iz, t, \epsilon) y = 0. \quad (8)$$

In an expression like  $e^{(1/\epsilon)f(t)}(e_0(t)\epsilon^k + e_1(t)\epsilon^{k+1} + \dots)$ , the term  $e_0(t)\epsilon^k$  shall be called the principal term.

We observe that the roots of the polynomial equation

$$w^2 + P_1(iz, t)w = 0 \quad (9)$$

are distinct except for a set of  $E^m$  measure zero. They are

$$w_1 = 0,$$

and

$$w_2 = -P_1(iz, t).$$

Hence by Lemma (1) we conclude the following:

$$|e^{\frac{1}{\epsilon}w_2}| < |e^{\frac{1}{\epsilon}w_1}| = 1. \quad (9')$$

LEMMA (2). For every value of  $i$  ( $i = 1, 2$ ) there exists an infinite number of functions  $u_{i0}(t), u_{i1}(t), \dots$ , continuous and with continuous derivatives of all orders such that  $u_{i0}(t)$  does not vanish at any point of  $J$  and if the functions

$$u_i(t, \epsilon) = \exp \left[ \frac{1}{\epsilon} \int_0^t w_i(s) ds \right] \sum_{j=0}^{m-1} u_{ij}(t) \epsilon^j$$

are substituted in the expression  $A(y)$  for  $y$ , then the coefficient of

$$\exp \left[ \frac{1}{\epsilon} \int_0^t w_i(s) ds \right] \epsilon^h \quad \begin{matrix} (i = 1, 2) \\ (h = 0, 1, 2, \dots, m) \end{matrix}$$

in the expansion thus obtained vanishes identically.

*Proof.* Write  $y(t, \epsilon) = \exp[1/\epsilon \int_0^t w(s) ds] v(t, \epsilon)$  where  $w(s)$  is some definite one of the two roots  $w_i(t)$ . We find the following:

$$\begin{aligned} y^{[1]}(t, \epsilon) &= \exp \left[ \frac{1}{\epsilon} \int_0^t w(s) ds \right] [w(t) v(t, \epsilon) + v^{[1]}(t, \epsilon)] \\ y^{[2]}(t, \epsilon) &= \exp \left[ \frac{1}{\epsilon} \int_0^t w(s) ds \right] ((w(t)^2 + \epsilon(d/dt) w(t)) v(t, \epsilon) \\ &\quad + 2w(t) v^{[1]}(t, \epsilon) + v^{[2]}(t, \epsilon)). \end{aligned}$$

Hence we conclude

$$A(y) = \exp \left[ \frac{1}{\epsilon} \int_0^t w(s) ds \right] \bar{A}(v) \quad (10)$$

where

$$\bar{A}(V(t, \epsilon)) = v^{[2]} + a_1(t, \epsilon) v^{[1]} + a_0(t, \epsilon) v \quad (11)$$

with  $a_1(t, \epsilon) = 2w(t) + P_1(iz, t)$ ,

$$a_0(t, \epsilon) = (w(t))^2 + \epsilon(d/dt) w(t) + P_1(iz, \epsilon) w(t) + P_2(iz, t, \epsilon).$$

For convenience we write:

$$\begin{aligned} a_2(t, \epsilon) &= 1, \\ a_n(t, \epsilon) &= 0. \quad (n > 2). \end{aligned}$$

If we let

$$a_n(t, \epsilon) \equiv \sum_{j=0}^{\infty} a_{nj}(t) \epsilon^j.$$

Then

$$a_{00}(t) = (w(t))^2 + P_1(iz, t) w(t) = 0 \quad (12)$$

$$a_{10}(t) = 2w(t) + P_1(iz, t) \quad (13)$$

so that  $a_{10}(t) \neq pmJ$ , the two roots  $w_i(t)$  being distinct. If we then insert in (10)

$$v(t, \epsilon) = \sum_{j=0}^{m-1} u_j(t) \epsilon^j,$$

we find the condition that the coefficient of

$$\exp \left[ \frac{1}{\epsilon} \int_0^t w(s) ds \right] \epsilon^h \quad (h = 0, 1, \dots, m)$$

in  $Ay(t, \epsilon)$  vanishes to be

$$\sum_{j+k+n=h} a_{ij}(t)(d^h/dt^h) y_k(t) = 0. \quad (14)$$

The equation (14) is true for  $h = 0$  by (12). For the case  $h \neq 0$  we can write (14) in the form:

$$\begin{aligned} a_{10}(t)(d/dt) u_{h-1}(t) + a_{01}(t) u_{h-1}(t) \\ + \sum_{\substack{k < h-1 \\ j+k+n=h}} a_{ij}(t)(d/dt)^h u_k(t) = 0. \end{aligned} \quad (15)$$

If in (14)  $k = h$ , then  $n = j = 0$  and the term corresponding to this set of values has a coefficient  $a_{00}(t) = 0$  if  $k = h - 1$ . Since this is the case, we have either  $n = 1, j = 0$  or  $n = 0, j = 1$ , and the corresponding terms are the first terms of equation (15).

It appears then that  $u_{h-1}(t)$  can be determined in term of  $u_{h-2}, u_{h-3}, \dots, u_0$  as a solution of certain linear differential equation of the first order which

has no singular points in  $J$ , since  $a_{10}(t) \neq 0$ . Thus  $u_0(t), u_1(t), \dots, u_{m-1}(t)$  can be obtained in succession from (14) for  $(h = 1, 2, \dots, m)$ . For each  $w_i(t)$  we obtain in this way a sequence of functions  $u_{i0}(t), u_{i1}(t), \dots$ , such that if the expressions

$$u_i(t, \epsilon) = \exp \left[ (1/\epsilon) \int_0^t w_i(s) ds \right] \sum_{j=0}^{m-1} u_{ij}(t) \epsilon^j$$

be substituted for  $y$  in  $\mathcal{A}(y)$  the coefficient of

$$\exp \left[ (1/\epsilon) \int_0^t w_i(s) ds \right] \epsilon^h \quad (i = 1, 2 \quad \text{and} \quad h = 0, 1, \dots, m)$$

vanishes by (15) since the conditions (14) are now satisfied for  $h = 1, 2, \dots, m$ . Furthermore the differential equation for  $u_{i0}(t)$  is homogeneous, so that, by taking for  $u_{i0}(t)$  a solution which is different from zero at one point of  $J$ , we are sure that  $u_{i0}(t)$  does not vanish at any point of  $J$ . Since the  $a_{ij}(t)$  were continuous,  $u_{i0}(t), u_{i1}(t), \dots$ , are also continuous with all of their derivatives. The sequence of functions  $u_{i0}(t), u_{i1}(t), \dots$ , has then the properties stated in the lemma.

LEMMA (3). *The homogeneous linear ordinary differential equation of order two with two solutions  $u_i(t, \epsilon)$  has the form*

$$B(y) = y^{[2]} + b_1(t, \epsilon) y^{[1]} + b_0(t, \epsilon) y = 0$$

for  $\epsilon \leq \epsilon_0 > 0$ .

$$|b_i(t, \epsilon)| \leq M : b_i(t, \epsilon) = \sum_{j=0}^{\infty} b_{ij}(t) \epsilon^j, \quad (i = 1, 2). \quad (16)$$

*The coefficients  $b_{ij}(t)$  which appears here are continuous with all their derivatives, and*

$$b_{10}(t) = P_1(iz, t) \quad \text{and} \quad b_{1j}(t) = 0 \quad (17)$$

*for all  $j > 0$ ; and if*

$$P_2(iz, t, \epsilon) = \sum_0^{\infty} P_{2j}(t) \epsilon^j,$$

*then*

$$b_{0j} = P_{2j}(t), \quad (j \leq m).$$

*Notice that  $b_{00}(t) = 0$ .*

*Letting,*

$$P_i(iz, t, \epsilon) = \sum_0^{\infty} P_{ij}(t) \epsilon^j,$$

then

$$b_{ij} = P_{ij}, \quad (i = 1, 2).$$

*Proof of Lemma (3).* The homogeneous linear differential equation of order two with solutions  $u_i(t, \epsilon)$  is

$$\begin{vmatrix} y^{[2]} & y^{[1]} & y \\ u_1^{[2]}(t, \epsilon) & u_1^{[1]}(t, \epsilon) & u_1(t, \epsilon) \\ u_2^{[2]}(t, \epsilon) & u_2^{[1]}(t, \epsilon) & u_2(t, \epsilon) \end{vmatrix} = 0. \quad (18)$$

For the element of this determinant we have

$$u_i^{[j]}(t, \epsilon) = \exp \left[ (1/\epsilon) \int_0^t w_i(s) ds \right] \sum_{k=0}^{m+j-1} \lambda_{ijk}(t) \epsilon^k \quad (19)$$

where,

$$\lambda_{ij0}(t) = [w_i(t)]^j u_{i0}(t). \quad (20)$$

Thus if we factor out of (18)

$$\prod_{i=1}^2 \exp \left[ (1/\epsilon) \int_0^t w_i(s) ds \right]$$

the differential equation takes the form

$$\beta_2(t, \epsilon) y^{[2]} + \beta_1(t, \epsilon) y^{[1]} + \beta_0(t, \epsilon) y = 0 \quad (21)$$

where  $\beta_i(t, \epsilon)$  are polynomials in  $\epsilon$ . We have for the principal term of  $\beta_2(t, \epsilon)$

$$\beta_{20}(t) = \begin{vmatrix} \lambda_{110} & \lambda_{100} \\ \lambda_{210} & \lambda_{200} \end{vmatrix}.$$

In view of (20) this last determinant may be written

$$\prod_{i=1}^2 u_{i0}(t) \begin{vmatrix} w_1(t) & 1 \\ w_2(t) & 1 \end{vmatrix}.$$

This, except for a set of  $E^m$  measure zero, is not zero at any point on  $J$  as the  $w_i(t)$  are distinct and the  $u_{i0}(t) \neq 0$  by Lemma (2). Therefore for  $\epsilon \leq \epsilon_0$

$$|\beta_i(t, \epsilon)/\beta_2(t, \epsilon)| \leq M : b_i(t, \epsilon) = |\beta_i(t, \epsilon)/\beta_2(t, \epsilon)| = \sum_{j=0}^{\infty} b_{ij}(t) \epsilon^j.$$

But we can write (21) in the form

$$y^{[2]} + b_1(t, \epsilon) y^{[1]} + b_0(t, \epsilon) y = 0. \quad (22)$$

Now the functions  $b_{ij}(t)$  and their derivatives are continuous since the  $w_i(t)$  and  $u_i(t)$  are of this character. The first part of the lemma is thus proved.

Now let  $j_0$  be the smallest value of  $j$  which, for some  $i$  and  $t$ ,

$$b_{ij}(t) \neq P_{ij}(t)$$

From (19) and (20) we see that the principal term of

$$B[u_i(t, \epsilon) - A(u_i(t, \epsilon))] = \overline{B - A}[u_i(t, \epsilon)] \quad (23)$$

is

$$\epsilon^{j_0} \left[ \sum_{k=0}^1 [b_{kj_0}(t) - P_{kj_0}(t)][w_i(t)]^k \right] u_{i0}(t). \quad (24)$$

Assume  $j_0 \leq m$  if possible. In each part of the difference (23) the coefficient of

$$\exp \left[ (1/\epsilon) \int_0^t w_i(s) ds \right] \epsilon^{j_0}$$

must then vanish, in the first since  $u_i(t, \epsilon)$  are solutions of  $B(y) = 0$ ; in the second part, by Lemma (2). Therefore from (24)

$$\sum_{k=0}^1 [b_{kj_0}(t) - P_{kj_0}(t)][w_i(t)]^k = 0 \quad (i = 1, 2)$$

we conclude that

$$b_{kj_0}(t) = P_{kj_0}(t), \quad (k = 0, 1, 2).$$

The  $w_i(t)$  being distinct. This is a contradiction. Hence  $j_0 > m$ . The proof of Lemma (3) is completed.

Consider the two equations

$$u_{tt} + P_1(t) u_t + (1/\epsilon) b_0 u = 0, \quad (25)$$

$$\begin{aligned} y_{tt} + P_1(t) y_t + (1/\epsilon) b_0 y &= [(1/\epsilon) b_0 - P_2] y \\ &= \epsilon^{m+1} Q(t, \epsilon) y \end{aligned} \quad (26)$$

where  $Q(t, \epsilon)$  is continuous in  $t$  and  $\epsilon$ .

Let  $y = u(t, \epsilon) v(t, \epsilon)$  with  $v(0, \epsilon) = 1$  and  $v_t(0, \epsilon) = 0$  and  $u(t, \epsilon) = u_i(t, \epsilon)$ ,  $i = 1, 2$ , given in Lemma (2). Substituting into (26) yields

$$\epsilon v_{tt} + (P_1 + 2\epsilon u_t/u) v_t = \epsilon^m Q v$$



which can be integrated to give

$$v(t) = 1 + \epsilon^m \int_0^t [u(s)]^{-2} \int_0^s Q(\sigma) v(\sigma) u^2(\sigma) \exp \left[ -\frac{1}{\epsilon} \int_\sigma^s P_1(l) dl \right] d\sigma ds.$$

Upon multiplying by  $u(t)$  and interchanging the order of integrations, the last equation becomes

$$y(t) = u(t) + \epsilon^m \int_0^t y(\sigma) K(t, \sigma, \epsilon) d\sigma \quad (27)$$

where

$$K(t, \sigma, \epsilon) = Q(\sigma) \int_\sigma^t [u(s)]^{-2} u(t) u(\sigma) \exp \left[ -\frac{1}{\epsilon} \int_\sigma^s P_1(l) dl \right] ds$$

which implies that  $K(t, \sigma, \epsilon)$  is bounded for  $0 < \epsilon \leq \epsilon_0$  and for both  $u = u_1$  or  $u = u_2$ .

The method of standard successive iterations shows that there exists a unique solution  $y(t)$  to equation (27). Furthermore, the solution  $y(t)$  is bounded for  $t$  in  $J$  and  $0 \leq \epsilon \leq \epsilon_0$ . As a consequence, we have

$$y_i(t) = u_i(t, \epsilon) + \epsilon^m E_0, \quad \text{and} \quad \frac{\delta y_i}{\delta t} = \frac{\delta u_i}{\delta t} + \epsilon^{m-1} E_1$$

where  $E_0$  and  $E_1$  are functions in  $\epsilon$  and others but bounded for  $0 \leq \epsilon \leq \epsilon_0$ .  $E_0$  can be derived from equation (27) and  $E_1$  can be found by taking the derivative of equation (27). We can sum the above results into the following theorem.

**THEOREM (1).** *The ordinary differential equation (8) has two independent solutions  $y_i(t, z, \epsilon)$  ( $i = 1, 2$ ) such that if the integer  $m$  is chosen at pleasure, then*

$$y_i(t, z, \epsilon) = u_i(t, \epsilon) + \epsilon^m E_0, \quad \text{and} \quad \frac{\delta y_i}{\delta t} = \frac{\delta u_i(t, \epsilon)}{\delta t} + \epsilon^{m-1} E_1.$$

Now let  $y_1$  and  $y_2$  be as in Theorem (1). Therefore the general solutions of equation (2) are:

$$Y_1(t, z, \epsilon) = A_1 y_1 + A_2 y_2, \quad Y_2(t, z, \epsilon) = B_1 y_1 + B_2 y_2.$$

Using the initial conditions (3) we compute the constants  $A_i$  and  $B_i$  as:

$$\begin{aligned} A_1 &= -(d/dt)(y_2(0, z, \epsilon))/H(z, \epsilon), & A_2 &= (d/dt)(y_1(0, z, \epsilon))/H(z, \epsilon) \\ B_1 &= (y_2(0, z, \epsilon))/H(z, \epsilon), & B_2 &= (-y_1(0, z, \epsilon))/H(z, \epsilon) \end{aligned} \quad (28)$$

where

$$H(z, \epsilon) = y_2(\epsilon)(d/dt) y_1(0, z\epsilon) - y_1(0, z, \epsilon)(d/dt) y_2(0, z, \epsilon).$$

By the fact that  $w_1(t) = 0$ ,  $w_2(t) = -P_1(iz, t)$  and by Theorem (1) we have

$$y_1(t, z, \epsilon) = \sum_{j=0}^{m-1} u_{1j}(t)\epsilon^j + \epsilon^m E_0 \quad (29)$$

$$y_2(t, z, \epsilon) = \sum_{j=0}^{m-1} u_{2j}(t)\epsilon^j \exp \left[ \frac{-1}{\epsilon} \int_0^t P_1(iz, is) ds \right] + \epsilon^m E_0.$$

Lemma (2) indicates that for all  $t$  in  $J$  none of  $u_{10}(t)$  and  $u_{20}(t)$  is equal to zero. Therefore, except for a set of  $E^m$  measure zero,  $H(z, \epsilon) \neq 0$  for all small  $\epsilon \geq 0$ . Consequently we have  $B_1(z, \epsilon)$ ,  $B_2(z, \epsilon)$ , and  $A_2(z, \epsilon)$  converging to zero as  $\epsilon$  tends to zero ( $B_1$ ,  $B_2$ ,  $A_2$  are converging as functions of  $\epsilon$ ; i.e.  $z$  is fixed.) Theorem (1) indicates that  $y_i(t, z\epsilon)$  are bounded ( $i = 1, 2$ ) in  $\epsilon$ . Therefore we conclude that, except for a set of  $E^m$  measure zero,  $Y_2(t, z, \epsilon)$  tends to zero as  $\epsilon$  tends to zero for all  $t$  in  $J$  and each  $z$  in  $E^m$ , while  $Y_1(t, z, \epsilon)$  goes to  $A_1(z, 0) y_1(t, z, 0)$ .

From (28) and (29) we can write

$$Y_1(t, z, \epsilon) = (u_{10}(t))/(u_{10}(0)) + o(\epsilon). \quad (30)$$

From equation (15) we can determine  $u_{10}(t)$  as the solution of the equation

$$P_1(iz, t) \frac{du}{dt} + P_2(iz, t, 0) = 0. \quad (31)$$

Now (30) and (31) together show that

$$Y_1(iz, t, \epsilon) \rightarrow Y_1(iz, t, 0) \quad \text{as } \epsilon \rightarrow 0$$

for all  $t$  in  $J$ , where  $Y_1(t, z, 0)$  is the solution of the equation

$$P_1(iz, t)(d/dt) y + P_2(iz, t, 0) = 0 \quad (32)$$

with the initial condition  $Y(0, z, 0) = 1$ . Now we sum the above results in the following lemma.

LEMMA (4). *If  $Y_j(t, z, \epsilon)$  are the solutions of equation (6) with the initial conditions (5) and satisfy the condition (7), then, except for a set of  $E^m$  measure zero, and for all  $z$  in  $E^m$  we have*

$$Y_1(t, z, \epsilon) \rightarrow Y_1(t, z, 0)$$

and

$$Y_2(t, z, \epsilon) \rightarrow 0$$

as  $\epsilon$  tends to zero and for all  $t$  in  $J$ .

## III

In this section we shall prove the main theorem of the paper.

**THEOREM (2).** *Let the degree of  $P_j(iz, \epsilon, t)$  in  $z$  be at most  $k$  where  $(j = 1, 2)$  and for all  $t$  in  $J$  assume that  $P_1(iz, t) \not\equiv 0$ . Let also  $u_0(t, x)$  be  $H_{p+k, x}$ , the continuously differentiable solution of the equation (1<sup>0</sup>). Suppose that  $Y_1(t, z, \epsilon)$  and  $Y_2(t, z, \epsilon)$  satisfy condition (7) in Lemma (1). Then equation (1) is an  $H_p$  stable equation.*

**Remark (1).** If neither  $P_1(iz, t)$  nor  $P_2(iz, t, \epsilon)$  depends on  $t$ , then we can prove that condition (7) is equivalent to another condition on the roots of the polynomial equation

$$\epsilon w^2 + P_1(iz)w + P_2(iz, \epsilon) = 0.$$

Those who are interested may see a special case in [3].

**Remark (2).** Notice that if  $P_1(iz, t) = 0$ , we will lose both solutions of equation (6) as well as the initial conditions (5) by passing to the limit; so the condition that  $P_1(iz, t)$  is not zero becomes necessary.

*Proof of Theorem (2).* As in Theorem (2) in [1], we can write

$$\begin{aligned} u(t, x, \epsilon) - u(t, x, 0) &= (1/\sqrt{2\pi^m}) \int_{E^m} e^{ix \cdot z} (Y_1(t, z, \epsilon) \hat{u}(0, z, \epsilon) \\ &- Y_1(t, z, 0) \hat{u}(0, z, 0)) dz + (1/\sqrt{2\pi^m}) \int_{E^m} e^{ix \cdot z} Y_2(t, z, \epsilon) \delta_t \hat{u}(0, z, \epsilon) dz. \end{aligned}$$

Using Fourier transform notations, we have

$$\begin{aligned} u(t, x, \epsilon) - u(t, x, 0) &= \mathcal{F}^{-1}(Y_1(t, z, \epsilon) \hat{u}(0, z, \epsilon) - Y_1(t, z, 0) \hat{u}(0, z, 0)) \\ &+ \mathcal{F}^{-1}(Y_2(t, z, \epsilon) \delta_t \hat{u}(0, z, \epsilon)). \end{aligned}$$

Hence

$$\begin{aligned} \|u(t, x, \epsilon) - u(t, x, 0)\|_p &= \left( \int_{E^m} |(1 + |z|^2)^{p/2} (\hat{u}(t, z, \epsilon) - \hat{u}(t, z, 0))|^2 dz \right)^{1/2} \\ &\leq \left( \int_{E^m} |(1 + |z|^2)^{p/2} (Y_1(t, z, \epsilon) \hat{u}(0, z, \epsilon) - Y_1(t, z, 0) \hat{u}(0, z, 0))|^2 dz \right)^{1/2} \\ &\quad + \left( \int_{E^m} |(1 + |z|^2)^{p/2} Y_2(t, z, \epsilon) \delta_t \hat{u}(0, z, \epsilon)|^2 dz \right) \\ &\leq \left( \int_{E^m} |(1 + |z|^2)^{p/2} ((Y_1(t, z, \epsilon) \hat{u}(0, z, \epsilon) - \hat{u}(0, z, 0)) \right. \\ &\quad \left. + (Y_1(t, z, \epsilon) - Y_1(t, z, 0)) \hat{u}(0, z, 0)|^2 dz \right)^{1/2} \\ &\quad + \left( \int_{E^m} |(1 + |z|^2)^{p/2} Y_2(t, z, \epsilon) \delta_t \hat{u}(0, z, \epsilon)|^2 dz \right)^{1/2} \end{aligned}$$

$$\begin{aligned} &\leq \left( \int_{E^m} |(1 + |z|^2)^{p/2} (Y_1(t, z, \epsilon) [\hat{u}(0, z, \epsilon) - \hat{u}(0, z, 0)])|^2 dz \right)^{1/2} \\ &\quad + \left( \int_{E^m} |(1 + |z|^2)^{p/2} (Y_1(t, z, \epsilon) - Y_1(t, z, 0)) \hat{u}(0, z, 0)|^2 dz \right)^{1/2} \\ &\quad + \left( \int_{E^m} |(1 + |z|^2)^{p/2} Y_2(t, z, \epsilon) \delta_t \hat{u}(0, z, \epsilon)|^2 dz \right)^{1/2}. \end{aligned}$$

By using condition (7), we get

$$\begin{aligned} &\int_{E^m} |(1 + |z|^2)^{p/2} Y_1(t, z, \epsilon) [\hat{u}(0, z, \epsilon) - \hat{u}(0, z, 0)]|^2 dz \\ &\leq C \int_{E^m} |(1 + |z|^2)^{p/2} (\hat{u}(0, z, \epsilon) - \hat{u}(0, z, 0))|^2 dz \end{aligned}$$

and by condition (3) it follows that

$$\int_{E^m} |(1 + |z|^2)^{p/2} (\hat{u}(0, z, \epsilon) - \hat{u}(0, z, 0))|^2 dz \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0 \quad (33)$$

for all  $t$  in  $J$ .

Lemma (4) together with condition (7) give the boundedness of  $Y_1(t, z, 0)$ ; hence, by the Lebesgue dominated convergence theorem, we have

$$\int_{E^m} |(1 + |z|^2)^{p/2} (Y_1(t, z, \epsilon) - Y_1(t, z, 0)) \hat{u}(0, z, 0)|^2 dz \rightarrow 0 \quad (34)$$

as  $\epsilon \rightarrow 0$  for all  $t$  in  $J$ .

Again conditions (4) and (7) and Lemma (4) give with the dominated convergence theorem

$$\int_{E^m} |(1 + |z|^2)^{p/2} Y_2(t, z, \epsilon) \delta_t \hat{u}(0, z, \epsilon)|^2 dz \rightarrow 0 \quad (35)$$

as  $\epsilon \rightarrow 0$ , for all  $t$  in  $J$ .

Hence (33), (34), and (35) imply the following:

$$\|u(t, x, \epsilon) - u(t, x, 0)\|_p \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0, \quad \text{for all } t \text{ in } J.$$

Theorem (2) is proved.

#### REFERENCES

1. NUR, H. S., Singular perturbation on linear partial differential equation with constant coefficients. Ph. D. thesis, Univ. Calif. Berkeley, 1967.
2. BIRKHOFF, G. D., On asymptotic character of the solutions of certain linear differential equations containing a parameter. *Trans. Am. Math. Soc.* 9, 219-231.
3. KUMANO-GO, HITOSHI, On singular perturbation of linear partial differential equations with constant coefficients. *Japan Acad. Proc.* (1959), 541-546.